



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for Histidine tag AND recombinant protein AND fu [Preview](#) [Go](#) [Clear](#)

Limits Preview/Index History Clipboard Details

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

- Search History will be lost after eight hours of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.
- Click on query # to add to strategy

| Search              | Most Recent Queries                                                                     | Time     | Result                |
|---------------------|-----------------------------------------------------------------------------------------|----------|-----------------------|
| <a href="#">#19</a> | Search <b>Histidine tag AND recombinant protein AND fusion protein</b>                  | 14:26:05 | <a href="#">99</a>    |
| <a href="#">#13</a> | Search <b>methods in enzymology[Journal] AND recombinant protein AND fusion protein</b> | 14:02:12 | <a href="#">60</a>    |
| <a href="#">#12</a> | Search <b>methods in enzymology[Journal] AND recombinant protein</b>                    | 13:57:35 | <a href="#">1263</a>  |
| <a href="#">#11</a> | Search <b>methods in enzymology[Journal]</b>                                            | 13:57:19 | <a href="#">14882</a> |
| <a href="#">#9</a>  | <b>Related Articles for PubMed (Select 10069435)</b>                                    | 13:43:18 | <a href="#">134</a>   |
| <a href="#">#6</a>  | Search <b>recombinant protein production AND tag AND review[Publication Type]</b>       | 13:38:03 | <a href="#">6</a>     |
| <a href="#">#5</a>  | Search <b>recombinant protein production AND tag</b>                                    | 13:37:46 | <a href="#">257</a>   |
| <a href="#">#3</a>  | Search <b>gst AND recombinant protein production AND MBP and histidine</b>              | 13:26:33 | <a href="#">1</a>     |
| <a href="#">#1</a>  | Search <b>gst</b>                                                                       | 13:25:28 | <a href="#">8279</a>  |

[Clear History](#)

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Aug 30 2004 06:52:01



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [Go] [Clear]  
Limits Preview/Index History Clipboard Details

About Entrez

[Display] Abstract Show: 20 Sort Send to Text

Text Version

☐ 1: Biotechnology (N Y). 1993 Oct;11(10):1138-43.

[Related Articles, Links](#)

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

## Use of peptide libraries to map the substrate specificity of a peptide-modifying enzyme: a 13 residue consensus peptide specifies biotinylation in *Escherichia coli*.

Schatz PJ.

Affymax Research Institute, Palo Alto, CA 94304.

I describe a technique for screening peptide libraries of over 10<sup>9</sup> independent clones for substrates of peptide-modifying enzymes. The peptides, linked to their genetic material by the lac repressor, are exposed to the enzyme and then screened by affinity purification on a receptor specific for the modified product. The enzyme characterized, *E. coli* biotin holoenzyme synthetase, normally adds biotin to a specific lysine residue in complex protein domains. The 13 residue substrate identified by this library screening approach is much smaller than the 75 amino acid required sequence of the natural substrate, and can function at either end of a fusion protein. The sequence is quite distinct at some positions from that region of the natural substrates, presumably because the peptides have to mimic the folded structure formed by the natural substrate. This technique should be useful for mapping the substrate specificity of a variety of peptide-modifying enzymes. In addition, small peptide substrates that are enzymatically biotinylated at a single site should be useful for a variety of purposes in labeling, purification, detection, and immobilization of proteins.

PMID: 7764094 [PubMed - indexed for MEDLINE]

[Display] Abstract Show: 20 Sort Send to Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Aug 30 2004 06:52:01